

Pittsburg State University

Pittsburg State University Digital Commons

Ichthyology Collection

Biology

2018

Pittsburg State University Ichthyology Field Notebooks

Karisa Boyer

Pittsburg State University, karisaboyer@joplinschools.org

Follow this and additional works at: <https://digitalcommons.pittstate.edu/ichthyology>



Part of the [Marine Biology Commons](#), and the [Zoology Commons](#)

Recommended Citation

Boyer, Karisa, "Pittsburg State University Ichthyology Field Notebooks" (2018). *Ichthyology Collection*. 1.
<https://digitalcommons.pittstate.edu/ichthyology/1>

This Article is brought to you for free and open access by the Biology at Pittsburg State University Digital Commons. It has been accepted for inclusion in Ichthyology Collection by an authorized administrator of Pittsburg State University Digital Commons. For more information, please contact digitalcommons@pittstate.edu.

Gash

1966

A LIST OF FISHES KNOWN FROM
SOUTHEASTERN KANSAS AND ENVIRONS

ORDER PETROMYZONTIFORMES

Family Petromyzontidae

Ichthyomyzon castaneus Girard-- Neosho River tributaries

ORDER AMIIFORMES

Family Amiidae

Amia calva
~~Amia nana~~ (Linnaeus) Neosho-Spring River system

ORDER LEPISTOSTEIFORMES

Family Lepisosteidae

Lepisosteus platostomus (Rafinesque) most of eastern Kansas
Lepisosteus osseus (Linnaeus) most of eastern Kansas.
Lepisosteus oculatus (Rafinesque) Neosho River
SPOTTED
ORDER CLUPEIFORMES

Suborder Clupeoidei

Family Clupeidae

Dorosoma cepedianum (LeSueur) most of Kansas

Suborder Salmonoidei

Family Salmonidae

Salmo gairdnerii (Richardson) "escapes" in Shoal Creek, planted elsewhere

Suborder Notopteroidei

Family Hiodontidae

Hiodon alosoides (Rafinesque) larger streams of eastern Kansas

ORDER CYPRINIFORMES

Suborder Cyprinoidei

Family Catostomidae

Cycaleptus elongatus (LeSueur) Neosho River
Ictiobus cyprinella (Valenciennes) eastern Kansas
Ictiobus niger (Rafinesque) eastern Kansas
Ictiobus bubalus (Rafinesque) eastern Kansas

SAVING THE KANSAS AND THE DOG

THE DOG

Family: *Canidae*

Canis latrans (Linnaeus) - *Canis latrans*

ORDER: *Carnivora*

Family: *Canidae*

Canis latrans (Linnaeus) - *Canis latrans*

ORDER: *Carnivora*

Family: *Canidae*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Canis latrans (Linnaeus) - *Canis latrans*

Carpoides carpio (Rafinesque) all of Kansas
Carpoides velifer (Rafinesque) Neosho River
Moxostoma carinatum (Cope) Spring River-Shoal Creek drainage (uncommon)
Moxostoma aureolum Shoal Creek
Moxostoma macrolepidotum piscilabrum (Trautman) Spring River-Neosho River
Moxostoma duquesnii (LeSueur) Spring River-Shoal Creek
Moxostoma erythrum (Rafinesque) eastern Kansas
Minytrema melanops (Rafinesque) southeastern and mideastern Kansas
Catostomus commersoni (Lacepede) Shoal Creek
Hypentelium nigricans (LeSueur) Spring River-Neosho River

Family Cyprinidae

Cyprinus carpio (Linnaeus) all of Kansas
Carassius auratus (Linnaeus) locally introduced
Notemigonus crysoleucas (Mitchill) all of Kansas
Semotilus atromaculatus (Mitchill) Spring River-Neosho River
Chrosomus erythrogaster (Rafinesque) Spring River, Neosho River
Hybopsis biguttata (Kirtland) eastern Kansas
Hybopsis amblops (Rafinesque) Spring River, Neosho River
Hybopsis storeriana (Kirtland) eastern Kansas
Hybopsis x-punctata (Hubbs and Crowe) Neosho River, Spring River
Phenacobius mirabilis (Girard) all of Kansas
Notropis parvobromus (Cope) most of Kansas
Notropis rubellus (Agassiz) eastern Kansas
Notropis umbratilis (Girard) eastern Kansas
Notropis chrysocephalus (Rafinesque) (N. cornutus) Spring River, uncommon
Notropis pilsbryi (Fowler) Spring River, eastern tributaries of Neosho River (N. zomale)
Notropis boops (Gilbert) southeastern Kansas
Notropis camurus (Jordan and Meek) southeastern Kansas
Notropis spilopterus (Cope) Neosho and Spring Rivers
Notropis lutrensis (Baird and Girard) all of Kansas
Notropis topeka (Gilbert) Dry Wood Creek
Notropis volucellus (Cope) southeastern Kansas
Notropis bethanani (Meek) eastern Kansas
Notropis stramineus (Cope) Spring and Neosho Rivers
Hybognathus placita (Girard) uncommon in Spring-Neosho River drainage
Dionda nubila (Forbes) Spring River, Shoal Creek
Pimephales tenellus (Girard) southeastern Kansas
Pimephales vigilax (Baird and Girard) southeastern Kansas
Pimephales notatus (Rafinesque) most of eastern Kansas
Pimephales promelas (Rafinesque) most of Kansas
Campestris anomalum (Rafinesque) all of Kansas

NOTROPIS CORNUTUS - SHOAL CK.
 1 black
 1 ventral bar

Suborder Siluridae

FAMILY: ICTALURIDAE

Ictalurus punctatus (Rafinesque) all of Kansas
Ictalurus punctatus natalis (LeSueur) eastern Kansas
Ictalurus melas (Rafinesque) all of Kansas
Ictalurus nebulosus (LeSueur) introduced, uncommon in Shoal Creek
Pylodictis olivaris (Rafinesque) most of Kansas
Noturus flavus (Rafinesque) eastern Kansas
Noturus nocturnus (Jordan and Gilbert) southeastern Kansas
Noturus exilis (Nelson) Spring and Neosho River drainages
Noturus miurus (Jordan) Spring and Neosho Rivers

ORDER ANGUILLIFORMES

Family Anguillidae

Anguilla rostrata (LeSueur) Neosho and Spring Rivers

ORDER CYPRINODONTIFORMES

Suborder Cyprinodontoides

Family Cyprinodontidae

Fundulus notatus (Rafinesque) eastern Kansas

Fundulus sciadicus (Cope) not yet taken in Kansas; Shoal Creek, Missouri

Fundulus catenatus (Storer) Shoal Creek, south of Joplin, Mo.; Dry Wood Creek, Max

Family Poeciliidae

Gambusia affinis (Baird and Girard) all of eastern and southeastern Kansas

ORDER MUGILIFORMES

Family Atherinidae

Labidesthes sicculus (Cope) all of southern Kansas

ORDER PERCIFORMES

Suborder Percoides

Family Serranidae

Roccus chrysops (Rafinesque) Spring and Neosho Rivers

Family Centrarchidae

Micropterus dolomieu (Lacepede) Spring and Neosho River systems

Micropterus punctulatus (Rafinesque) all of southern Kansas

Micropterus salmoides (Lacepede) all of Kansas

Chaenobryttus glaucus (Cuvier) eastern Kansas

Lepomis cyanellus (Rafinesque) all of Kansas

Lepomis microlophus (Gunther) introduced, uncommon in Shoal Creek

Lepomis megalotis (Rafinesque) eastern Kansas

Lepomis humilis (Girard) all of Kansas

Lepomis macrochirus (Rafinesque) all of Kansas

Ambloplites rupestris (Rafinesque) Spring and Neosho Rivers

Pomoxis nigromaculatus (DeSueur) all of Kansas

Pomoxis annularis (Rafinesque) all of Kansas

Family Percidae

Stizostedion canadense (Smith) Neosho and Spring Rivers

Percina phoxocephala Nelson) southeastern Kansas

Percina caprodes (Rafinesque) eastern Kansas

Percina shumardi (Girard) Neosho and Spring Rivers

Percina copelandi (Jordan) southeastern Kansas

Etheostoma stigmaeum (Jordan) Spring and Neosho Rivers

Etheostoma nigrum - subsp. Spring River, Shoal Creek, Dry Wood Creek

Etheostoma chlorosomum (Hay) Neosho and Spring Rivers

Amphibia (Lacerta) Lacerta and Spring River

ORDER CYPRINODONTIFORMES

Suborder Cyprinodontiformes

Family Cyprinodontidae

Brachydanio rerio (Rafinesque) eastern Kansas

Brachydanio rerio (Günther) not yet taken in Kansas; Shoal Creek, Missouri
Brachydanio rerio (Günther) Shoal Creek, south of Topeka, Mo.; Dry Wood Creek, Tex.

Family Poeciliidae

Gambusia affinis (Baird and Girard) all of eastern and southern Kansas

ORDER MOLLUSCOFORMES

Family Acanthinoidae

Acanthinoides alatus (Günther) all of southern Kansas

ORDER REMIPEDIFORMES

Suborder Remipediformes

Family Serripidae

Remipes caryocarpus (Rafinesque) Spring and Neosho Rivers

Family Centrarchidae

Micropterus dolomieu (Lacépède) Spring and Neosho River systems

Micropterus dolomieu (Rafinesque) all of southern Kansas

Micropterus dolomieu (Lacépède) all of Kansas

Micropterus dolomieu (Günther) eastern Kansas

Micropterus dolomieu (Rafinesque) all of Kansas

Micropterus dolomieu (Günther) introduced, introduced to Shoal Creek

Micropterus dolomieu (Rafinesque) eastern Kansas

Micropterus dolomieu (Günther) all of Kansas

Micropterus dolomieu (Rafinesque) all of Kansas

Micropterus dolomieu (Rafinesque) Spring and Neosho Rivers

Micropterus dolomieu (Günther) all of Kansas

Micropterus dolomieu (Rafinesque) all of Kansas

Family Percidae

Micropterus dolomieu (Günther) Spring and Neosho Rivers

Micropterus dolomieu (Rafinesque) southern Kansas

Micropterus dolomieu (Lacépède) eastern Kansas

Micropterus dolomieu (Günther) Spring and Neosho Rivers

Micropterus dolomieu (Jordan) southeastern Kansas

Micropterus dolomieu (Jordan) Spring and Neosho Rivers

Micropterus dolomieu (Jordan) Spring and Neosho Rivers

Micropterus dolomieu (Jordan) Spring and Neosho Rivers

Etheostoma zonale (Cope) Spring and Neosho Rivers
Etheostoma blenniodes subsp. Spring River, Shoal Creek
Etheostoma whipplii (Girard) southeastern Kansas
Etheostoma punctulatum (Agassiz) Spring River, Shoal Creek
Etheostoma cragini (Gilbert) southeastern Kansas
~~Etheostoma~~ Etheostoma spectabile (Agassiz) most of Kansas
Etheostoma flabellare (Rafinesque) Spring and Neosho Rivers
Etheostoma gracile (Girard) Spring River drainage
Etheostoma microperca (Jordan and Gilbert) Shoal Creek, just inside Missouri

Family Sciaenidae

Aplodinotus grunniens (Rafinesque) most of Kansas

Suborder Cottoidei

Family Cottidae

Cottus caroliniae (Gill) Spring River And Neosho River

A LIST OF FISHES KNOWN FROM SOUTHEASTERN KANSAS AND ENVIRONS

- small stream fish soft integument.*
1. Ichthyomyzon castaneus Girard Chestnut lamprey-Neosho river tributaries
  2. Amia calva Linnaeus Bowfin-Neosho and Spring River systems
PRIMITIVE line form. Ganooid scales - homocercal tail
  3. Lepisosteus platostomus Rafinesque Shortnose gar-most of eastern Kansas
fin yellowish. spots on 50cm GAR 1.5cm. in DIA.
  4. Lepisosteus osseus (Linnaeus) Longnose gar - most of eastern Kansas
Spanning in later May-June. BEAK 2-2.5X that of shortnose
  5. Lepisosteus oculatus (Cope) Spotted gar - Neosho River system
first specimen (Spring) 1963. St. Paul Refuge.
  6. Dorosoma cepedianum (LeSueur) Gizzard shad - most of Kansas
perlen sense Threadfin chub
  7. Salmo gairdnerii Richardson Rainbow trout - escaped to Shoal Creek
soft rays, deep skin covered scales
  8. Hiodon alosoides (Rafinesque) Goldeye - larger streams and lakes
OSAGE R. large teeth
  9. Cycoreus elongatus (LeSueur) Blue sucker - Neosho River system
PA. fin smallmouth Pony mouth
  10. Ictiobus cyprinella (Valenciennes) Bigmouth buffalofish - eastern Kansas
MAX - 20-30 lbs
  11. Ictiobus niger (Rafinesque) Black buffalofish - eastern Kansas
  12. Ictiobus bubalus (Rafinesque) Smallmouth buffalofish - eastern Kansas
hard to distinguish
  13. Carpionodes carpio (Rafinesque) River carpsucker-all of Kansas
of dorsal fin extended, snout at apex of lower lip. - 5 lbs max
  14. Carpionodes velifer (Rafinesque) Highfin carpsucker-Neosho River system
hump on anterior third of dorsal fin. coloration similar
  15. Moxostoma carinatum (Cope) River redhorse-Spring River/Shoal Creek
14-20 lbs 2 1/2 ft in length good eaters - breeding colors beautiful
  16. Moxostoma aureolum (LeSueur) Shorthead redhorse-Shoal Creek
  17. Moxostoma macrolepidotum pisolabrum (Trautman) Eastern redhorse-Spring and Neosho River systems
  18. Moxostoma duquesnii (LeSueur) Black redhorse-Spring River and Shoal Creek
  19. Moxostoma erythrurum (Rafinesque) Golden redhorse-eastern Kansas
  20. Mnytrema melanops (Rafinesque) Spotted sucker-eastern Kansas
well defined melanistic spots - mauling in bands
  21. Catostomus commersoni (Lacepede) White sucker-eastern Kansas
  22. Hypentelium nigricans (LeSueur) Northern hogsucker-Spring and Neosho River systems
  23. Cyprinus carpio Linnaeus All of Kansas
  24. Carassius auratus (Linnaeus) Goldfish-introduced
  25. Notemimonus crysoleucas (Mitchill) Golden shiner-all of Kansas
teeth 1-50
  26. Semotilus atromaculatus (Mitchill) Creek chub-Spring and Neosho Rivers Marmaton and Osage River systems
  27. Chrosomus erythrogaster (Rafinesque) Southern Redbelly Dace-Spring and Neosho River systems

GAR'S.

(3-5): SPAWNING: Typically for every ♀, there will be 2 ♂, swimming in tandem.

Spawning takes place in weeds. ♀ spin when releasing eggs. Eggs are pellets. **TOXIC!!! FATAL.**
Egg covers highly toxic.

8 - surface feeders - insectivorous; 8-10 ft deep.
Clear water - length to .5 meters, to 30-40

all suckers have cycloid scales

~~5~~ Calostomidae - no spine

- in dorsal fin 10+ rays.

inferior mouth, lips thickened

pharyngeal teeth 1 row typically 16+

Cyprinidae - at least 1 fin w/ ^{usually} spines
usually not more than 2 spines

dorsal fin 10 or less rays

lips typically thinner

mouth - not inferior - teeth 8 or less

~~Cyprinidae~~

28. Hybopsis biguttata (Kirtland) Hornyhead chub-eastern Kansas
29. Hybopsis amblops (Rafinesque) Bigeye chub-Spring and Neosho Rivers
30. Hybopsis storeriana (Kirtland) Silver chub-eastern Kansas
31. Hybopsis x-punctata Huobbs and Crowe Gravel chub-Neosho and Spring Rivers
32. Phenacobius mirabilis (Girard) Suckermouth minnow-all of Kansas
- X 33. Notropis percobromus (Cope) Plains shiner-most of Kansas
34. Notropis rubellus (Agassiz) Rosyface shiner-eastern Kansas
35. Notropis umbratilis (Girard) ^{scales deep-smooth} Redfin shiner-eastern Kansas
36. Notropis chrysocephalus (Rafinesque) (cornutus) Common shiner-Spring R.
37. Notropis pilsbryi Fowler Spring River; eastern tributaries of Neosho R.
38. Notropis boops Gilbert Bigeye shiner-southeastern Kansas ^{cold fast water - specific preferences}
39. Notropis camurus (Jordan and Meek) bluntface shiner-southeastern Kansas
40. Notropis spilopterus (Cope) Spotfin shiner-Neosho and Spring Rivers
41. Notropis lutrensis (Baird and Girard) Red shiner-all of Kansas ^{usually 20-30 cm deep still water}
- X 42. Notropis volucellus (Cope) Mimic shiner-southeastern Kansas
43. Notropis buchmanii Meek Ghost shiner-eastern Kansas
44. Notropis topeka Gilbert Topeka shiner-eastern Kansas
45. Notropis stramineus (Cope) Spring and Neosho Rivers
46. Hybognathus placita Girard Plains minnow-Spring and Neosho Rivers ^{usually moderately heavily pigmented}
47. Dionda nubila (Forbes) Ozark minnow-Spring River and Shoal Creek ^{8-10 cm distinct separation of pigment on lat. line.}
48. Pimephales tenellus (Girard) ^{resembles P. notatus in color, V-shaped caudal spot.} Slim minnow-southeastern Kansas
49. Pimephales vigilax (Baird and Girard) Bullhead minnow-southeastern Kansas
50. Pimephales notatus (Rafinesque) Bluntnose minnow-eastern Kansas ^{NO 3 DL 3-11-68}
51. Pimephales promelas Rafinesque Fathead minnow-most of Kansas
52. Campostoma anomalum (Rafinesque) Stoneroller ^{small scales, very smooth skin.} (rotgut) minnow-Kansas
53. Ictalurus punctatus (Rafinesque) Channel catfish-all of Kansas ^{MAX. 30 lbs (APPROX.)}
54. Ictalurus furcatus (LeSueur) Blue catfish-Neosho and Spring Rivers ^{HARD TO DISTING. hump on back of head - B.G.}
55. Ictalurus natalis (LeSueur) Yellow bullhead-all of Kansas
56. Ictalurus melas (Rafinesque) Black bullhead-all of Kansas

B.G.

Ictalurus - prominent adipose fin
dorsal ray - Bullheads A. 25+
other Ictalurids A. 19 or fewer

~~sub~~ flesh in Bullheads remains red after skinning.

DORSAL
VENTRAL



69. Roccus caryops - regius sandy bottom -
depth 6-20 ft. must be on some
kind of pennant.
strictly predatory.

mottled

57. Ictalurus nebulosus (LeSueur) Brown bullhead-Introduced; uncommon in Shoal Creek and in Lake Farlington
58. Pyiodictis olivaris (Rafinesque) Flathead catfish-widely distributed
(YELLOW CAT)... non-migrating head shape - - larger bodies of males 109/bs
59. Noturus flavus Rafinesque Stonecat-all of Kansas
60. Noturus nocturnus Jordan and Gilbert Freckled madtom-southeastern Kansas
61. Noturus exilis Nelson Slender madtom-Spring and Neosho River drainages
6 known protein toxins on spines - *3 Drywood 3-11-66
62. Noturus miurus Jordan Brindled madtom-Neosho and Spring Rivers
63. Anguilla rostrata (LeSueur) Neosho, Spring and Verdigris Rivers
SPAW AT SARGASSO SEA, BURMUDA. American eel
64. Fundulus catenatus (Storer) Northern studfish-Shoal Creek, south of Joplin Mo.; not yet taken from Kansas Drywood
WIDE HABITAT RANGE
65. Fundulus sciadicus (Cope) Plains topminnow-Shoal Creek east of state line; not taken from Kansas #3, 1965
66. Fundulus notatus (Rafinesque) Blackstripe topminnow-eastern Kansas - 3 Dry W. 7-11-66
67. Gambusia affinis (Baird and Girard) Mosquitofish-eastern and southern Kansas
(wide spread Genus) Kansas not in strip pits
68. Labidesthes sicculus (Cope) Brook silverside-all of southern Kansas #3. D.W. 3-11-66
69. Roccus chrysops (Rafinesque) White bass-Spring and Neosho Rivers
70. Micropterus dolomieu Lacepede Smallmouth bass-Spring and Neosho River systems
71. Micropterus punctulatus (Rafinesque) Spotted bass-southern Kansas
72. Micropterus salmoides (Lacepede) Largemouth bass-all of Kansas
73. Chaenobryttus gulosus (Cuvier) Warmouth sunfish-eastern Kansas
74. Lepomis cyanellus Rafinesque Green sunfish-all of Kansas
75. Lepomis microlophus (Gunther) Redear sunfish-introduced; Shoal Creek some strip pits and probably Lake Farlington
76. Lepomis megalotis (Rafinesque) Long-ear sunfish-eastern Kansas #3. Dry Wood 5-11-66
77. Lepomis humilis (Girard) Orange-spot sunfish-all of Kansas
78. Lepomis macrochirus Rafinesque Bluegill sunfish-all of Kansas
79. Ambloplites rupestris Rafinesque Rockbass-Spring and Neosho Rivers
East Cow Creek
80. Pomoxis nigromaculatus (LeSueur) Black crappie-all of Kansas
81. Pomoxis annularis Rafinesque White crappie-all of Kansas
82. Stizostedion canadense (Smith) Sauger-Neosho and Spring River drainages
83. Percina phoxocephala (Nelson) Slenderhead darter-southeastern Kansas
84. Percina caprodes (Rafinesque) Logperch-eastern Kansas #3 Dry Wood Mar. 11, 1966

PERCIDAE - no swim bladder - require -
high O₂ level - 6-8 ppm D.O₂

L. siculus ends at start of brackish
water (salinity).

Menidia audens -

highly saline waters
swim is shorter

- 4
85. Percina shumardi (Girard) River darter-Neosho and Spring River systems
  86. Percina copelandi (Jordan) Channel darter-southeastern Kansas
  87. Etheostoma stigmaeum (Jordan) Speckled darter-Spring and Neosho Rivers
ETHEOSTOMA - Regular lateral line system
  88. Etheostoma nigrum Rafinesque Johnny darter-Spring River, Shoal and Drywood Creeks (subsp.)
leaf buds in winter
#3 D.W. 3-11-66
  89. Etheostoma chlorosomum (Hay) Bluntnose darter-Neosho and Spring Rivers
  90. Etheostoma zonale (Cope) Banded darter-Spring and Neosho Rivers
  91. Etheostoma blennioides Rafinesque Spring River and Shoal Creek (subsp.)
  92. Etheostoma whipplei (Girard) Redfin darter-southeastern Kansas
  93. Etheostoma punctulatum (Agassiz) Stippled darter-Spring River and Shoal Creek
Chassid fish
  94. Etheostoma cragini Gilbert Arkansas darter-southeastern Kansas
  95. Etheostoma spectabile (Agassiz) Orangethroat darter-most of Kansas
#3 D.W. MAR. 11, 1966
  96. Etheostoma flabellare Rafinesque Fantail darter-Spring and Neosho Rivers
Drywood Creek
  97. Etheostoma gracile (Girard) Slough darter-Spring River drainage
  98. Etheostoma microperca Jordan and Gilbert Shoal Creek, immediately east of state line
  99. Aplodinotus grunniens Rafinesque Freshwater drum-most of Kansas
  100. Cottus carolinae (Gill) Banded sculpin-Spring and Neosho Rivers
+ SHOAL CREEK.

Branley A. Branson 1963

J. Carl Bass
KSCP - 1966

$$D.O_2 = \frac{1}{T}$$

BIOLOGY 333
ICHTHYOLOGY

Blair, F. W., A. P. Blair, P. Brodkorb, F. R. Cagle and G. A. Moore
VERTEBRATES OF THE UNITED STATES 1957; McGraw-Hill Co., New York.

Classification structure relating to fishes of North America and
specifically to Southeast Kansas:

Phylum CHORDATA

Subphylum CRANIATA (VERTEBRATA)

Superclass AGNATHA

Class Petromyzontes

p. 37

Order Petromyzontiformes

p. 37

Family Petromyzontidae

Ichthyomyzon castaneus chestnut lamprey p. 42

Superclass GNATHOSTOMATA

Class Teleostomi

Order Acipenseriformes

p. 51

Family Polydontidae paddlefish

Acipenseridae sturgeon

Order Amiiformes

p. 53

Family Amiidae bowfin

Order Lepisosteiformes

p. 54

Family Lepisosteidae gar-fishes

Order Clupeiformes

p. 56

Family Elopidae tarpon

Clupeidae herrings and shad

Engraulidae anchovies

Salmonidae salmon, trout, char

Coregonidae whitefish

Thymallidae grayling

Osmeridae smelt

Umbridae mudminnow

Esocidae pickerel, pike, muskellunge

Hiodontidae mooneye

Order Cypriniformes

p. 74

Family Catostomidae suckers

Cyprinidae minnows (p. 93)

Ictaluridae catfishes (p. 139)

Order Anguilliformes

p. 145

Family Anguillidae eels

Order Beloniformes needlefish (coastal)

p. 146

Family Belonidae needlefish

Order Gadiformes

p. 146

Family Gadidae burbot

Order Gasterosteiformes

p. 146

Family Gasterosteidae stickleback

Order Syngnathiformes

p. 147

Family Syngnathidae pipefishes

Order Cyprinodontiformes

p. 148

Family Amblyopsidae blindfishes

Cyprinodontidae killifish

Poeciliidae livebearers

Order Percopsiformes

p. 159

Family Percopsidae troutperch

Aphredoderidae pirateperch

Order Mugiliformes

p. 161

Family Atherinidae silversides

Mugilidae mullet

U.S. DEPARTMENT OF AGRICULTURE
BUREAU OF PLANT INDUSTRY

PLANT INDUSTRY DIVISION
WASHINGTON, D. C.

1950

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION
WASHINGTON, D. C.

1950

PLANT INDUSTRY DIVISION

1950

PLANT INDUSTRY DIVISION

1950

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

WASHINGTON, D. C.

PLANT INDUSTRY DIVISION

Order Perdiiformes

p. 162

Family Centropomidae snook (coastal)
Serranidae basses
Centrarchidae sunfishes
Percidae perch, darters
Sciaenidae drum
Sparidae porgy, sheephead
Cichlidae cichlids (Rio Grande perch)
Embiotocidae surfperch
Eleotridae sleeper (coastal)
Gobiidae goby
Cottidae sculpins

Order Pleuronectiformes

p. 205

Family Pleuronectidae flounder (coastal)
Achiridae sole (coastal)

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966


Area Second Cow Creek Station No. 1 Date Feb. 24, 1966
 Location Sec. 9, T. 30 S., R. 23 E., Crawford Co., Kansas
 Water Temp.(F.) 39° Air Temp.(F.) 48° Turbidity(ft.) 5-6
 Rate of Flow(cu. ft./min.) 3-4 Type of Drainage grassland, some trees
 Habitat rock bottom, heavy leaf cover, little aquatic plant growth
 Specimens Taken

ORDER	FAMILY	Species	Remarks
CYPRINIFORMES	Cyprinidae	<u>Phenacobius mirabilis</u> <u>Notropis umbratilis</u> <u>Pimephales notatus</u>	
CYPRINODONTIFORMES	Cyprinodontidae Poeciliidae	<u>Fundulus notatus</u> <u>Gambusia affinis</u>	
MUGILIFORMES	Atherinidae	<u>Labidesthes sicculus</u>	
PERCIFORMES	Centrarchidae	<u>Lepomis cyanellus</u>	

COLLECTION DATA
 DATE: 1968
 LOCATION: Second Cow Creek
 T. 30 S., R. 23 E., Crawford Co., Kansas
 DATE OF FLOW (or, if (min.) 3-4 Type of drainage: grassland, some trees
 WATER TEMP. (°F.) 39° AIR TEMP. (°F.) 48°
 WIND DIRECTION: S-S
 WIND VELOCITY: 0-5
 MOON: 1st Quarter
 TIME OF DAY: 10:00 AM
 COLLECTOR: J. H. ...
 SPECIES TAKEN:

ORDER	FAMILY	SPECIES	REMARKS
CYPRINIDAE	Cyprinidae	<u>Phenacodus nigricollis</u> <u>Notropis rubrivittis</u> <u>Phoxinotus notatus</u>	
CYPRINODONTIDAE	Cyprinodontidae	<u>Gambusia notata</u> <u>Gambusia affinis</u>	
MISCELLANEOUS	Atherinidae	<u>Atherinops affinis</u>	
POCILICHTHIDAE	Pocilichthidae	<u>Pocilichthys cyanoptera</u>	

COW CREEK SURVEY
SPRING 1963
Dept. of Biology
Kansas State College
Pittsburg, Kansas



COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Dry Wood Creek Station No. 1 Date Mar. 3, 1966
 Location Sec. 7, T. 28 S., R. 23 E., Crawford Co., Kansas
 Water Temp.(F.) 48 Air Temp.(F.) 58 Turbidity(ft.) 4 ft.
 Rate of Flow(cu. ft./min.) 24 Type of Drainage grassy-wooded area
 Habitat rock bottom, much aquatic vegetation, especially Spirogyra; some organic
debris in riffles, i.e., sticks, etc.
 Specimens Taken

ORDER	FAMILY	Species	Remarks
CYPRINIFORMES	Cyprinidae	<u>Notemigonus crysoleucas</u> <u>Notropis lutrensis</u>	
	(Siluroidei)		
	Ictaluridae	<u>Ictalurus melas</u>	
CYPRINODONTIFORMES	Cyprinodontidae	<u>Fundulus notatus</u>	
MUGILIFORMES	Poeciliidae	<u>Labidesthes sicculus</u>	
PERCIFORMES	Centrarchidae	<u>Micropterus salmoides</u>	
		<u>Lepomis cyanellus</u>	
		<u>Lepomis megalotis</u>	
		<u>Lepomis macrochirus</u>	
		<u>Percina caprodes</u>	
	Percidae	<u>Etheostoma spectabile</u>	

1911

1950

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Dry Wood Creek Station No. 2 Date March 3, 1966
 Location Sec. 32, T. 27 S., R. 23 E., Crawford Co. Kansas
 Water Temp.(F.) 47° Air Temp.(F.) 59° Turbidity(ft.) 4-5 ft.
 Rate of Flow(cu. ft./min.) 30 Type of Drainage forest-grassland
 Habitat rock bottom, rapid current, few aquatic plants

Specimens Taken

ORDER	FAMILY	Species	Remarks
CYPRINIFORMES	Cyprinidae	<u>Semotilus atromaculatus</u> <u>Notropis umbratilis</u> <u>Pimephales notatus</u> <u>Campostoma anomalum</u>	
CYPRINODONTIFORMES	Cyprinodontidae Poeciliidae	<u>Fundulus notatus</u> <u>Gambusia affinis</u>	
PERCIFORMES	Centrarchidae Percidae	<u>Micropterus salmoides</u> <u>Etheostoma nigrum</u> <u>Etheostoma spectabile</u> <u>Etheostoma flabellare</u>	

COLLECTED DATA

ICHTHYOLOGICAL SURVEY, Spring, 1968

Station No. 2 Date March 2, 1968

Location Sec. 29, T. 29 N., R. 29 E., Crawford Co., Kansas

Water Temp. (F.) 59° Air Temp. (F.) 59° Turbidity (ft.) 1-2 ft.

Rate of Flow (cu. ft./min.) 30 Type of Drainage Forest-prairie

Habitat rock bottom, slight current, few aquatic plants

Specimens Taken

ORDER	FAMILY	SPECIES	REMARKS
CYPRINIDAE	Cyprinidae	<u>Lepomis microlophus</u>	
		<u>Notropis umbellatus</u>	
		<u>Limnodynastes notatus</u>	
		<u>Desmognathus fusca</u>	
CYPRINIDAE	Cyprinodontidae	<u>Leuciscus notatus</u>	
		<u>Carassius auratus</u>	
CYPRINIDAE	Cyprinodontidae	<u>Leuciscus notatus</u>	
		<u>Desmognathus fusca</u>	
		<u>Leuciscus notatus</u>	
		<u>Desmognathus fusca</u>	

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Dry Wood Creek Station No. 3 Date March 10, 1966Location Sec. 32, T. 27 S., R. 23 E., Crawford Co., KansasWater Temp.(F.) 59° Air Temp.(F.) 65° Turbidity(ft.) 4Rate of Flow(cu. ft./min.) 40 (approx.) Type of Drainage farmed grassland,Habitat mud bottom, few rocks; oak overhang provides leaf covering of bottom in
some spots

Specimens Taken

ORDER	FAMILY	Species	Remarks
CLUPEIFORMES	Clupeidae	<u>Dorsoma cepedianum</u>	
CYPRINIFORMES	Catostomidae	<u>Ictiobus bubalus</u>	
		<u>Catostomus commersonii</u>	
	Cyprinidae	<u>Cyprinus carpio</u>	
		<u>Notemigonus crysoleucas</u>	
		<u>Notropis umbratilis</u>	
		<u>Notropis lutrensis</u>	
		<u>Pimephales notatus</u>	
		<u>Camptostoma anomalum</u>	
	Ictaluridae	<u>Ictaluris natalis</u>	
		<u>Ictaluris melas</u>	
		<u>Noturus exilis</u>	
CYPRINODONTIFORMES	Cyprinodontidae	<u>Fundulus notatus</u>	
MUGILIFORMES	Atherinidae	<u>Labidesthes sicculus</u>	
PERCIFORMES	Percidae	<u>Micropterus salmoides</u>	
		<u>Lepomis cyanellus</u>	
		<u>Lepomis megalotus</u>	
		<u>Lepomis humilis</u>	
		<u>Lepomis macrochirus</u>	
		<u>Pomoxis nigromaculatus</u>	
		<u>Pomoxis annularis</u>	
		<u>Percina caprodes</u>	
		<u>Etheostoma nigrum</u>	
		<u>Etheostoma spectabile</u>	
		<u>Etheostoma flabelare</u>	

ICHTHYOLOGY 333, Spring, 1966

Location Sec. 30, T. 27 S., R. 24 E., Crawford Co., Kansas

Rate of Flow(cu. ft./min.) ? - swift Type of Drainage wooded hills- typical Ozarkian area

Habitat limestone bottom, heavy filamentous algae growth

Remarks

CYPRINIFORMES	Ictaluridae	<u>Ictaluris melas</u>	taken by R. Streeter

2001

2010

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Kansas State Quail Farm Station No. - Date March 16, 1966Location Sec. 23, T. 30 S., R. 24 E., Crawford Co., KansasWater Temp.(F.) 55° Air Temp.(F.) 73° Turbidity(ft.) 2-3Rate of Flow(cu. ft./min.) 0 Type of Drainage strip mine lakeHabitat shale bottom and bank, moderate vegetation in water; little overhang by trees in area, rapidly descending banks

Specimens Taken

ORDER	FAMILY	Species	Remarks
PERCIFORMES	Centrarchidae	<u>Micropterus salmoides</u>	SL: 87 mm, wt.: 12.7g hit 1" top-water plug time: 2-3: P.M. wind: 0-5 mph NNW partly cloudy
		<u>Chaenobryttus gluosus</u>	April 7, 1966

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Schemerhorn

Area Shoal Creek

Station No. Park Date March 17, 1966

Location Sec. 35, T. 34 S., R. 25 E., Cherokee Co., Kansas

Water Temp.(F.) 64° Air Temp.(F.) 75° Turbidity(ft.) 2-3

Rate of Flow(cu. ft./min.) very rapid Type of Drainage rocky wooded hill country

Habitat rock bottom, loose gravel-size particles; some large boulders

Specimens Taken

ORDER	FAMILY	Species	Remarks
CYPRINIFORMES	Catostomidae	<u>Moxostoma erythrum</u>	
		<u>Minytrema melanops</u>	
	Cyprinidae	<u>Semotilus atromaculatus</u>	
		<u>Hybopsis x-punctata</u>	
		<u>Notropis umbratilis</u>	
		<u>Notropis pilsbryi</u>	
		<u>Notropis lutrensis</u>	
		<u>Notropis boops</u>	
		<u>Pimephales notatus</u>	
		<u>Campostoma anomalum</u>	
CYPRINODONTIFORMES	Cyprinodontidae	<u>Fundulus notatus</u>	
	Poeciliidae	<u>Gambusia affinis</u>	
PERCIFORMES	Centrarchidae	<u>Micropterus ~~salmoides~~ punctulatus</u>	
		<u>Micropterus salmoides</u>	
		<u>Lepomis cyanellus</u>	
		<u>Lepomis megalotus</u>	
		<u>Lepomis humilis</u>	
		<u>Lepomis macrochirus</u>	
		<u>Percina caprodes</u>	
		<u>Etheostoma nigrum</u>	
	Percidae	<u>Etheostoma spectabile</u>	

1897-1898

Specimens Taken

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

State Line

Area Shoal Creek (1 M. west of St. line) Station No. Station Date March 19, 1966

Location Sec. 35, T. 34 S., R. 25 E., Cherokee Co., Kansas

Water Temp.(F.) 58° Air Temp.(F.) 75° Turbidity(ft.) 2-3

Rate of Flow(cu. ft./min.) swift Type of Drainage rock-oaken hills

Habitat limestone bottom, little vegetation in water

Specimens Taken

ORDER	FAMILY	Species	Remarks
CYPRINIFORMES	Cyprinidae	<u>Semotilus atromaculatus</u> <u>Chrosomus erythrogaster</u> <u>Hybopsis x-punctata</u> <u>Notropis umbratilis</u> <u>Notropis pilsbryi</u> <u>Notropis lutrensis</u> <u>Campostoma anomalum</u>	
CYPRINODONTIFORMES	Cyprinodontidae	<u>Fundulus notatus</u>	
PERCIFORMES	Percidae	<u>Etheostoma flabellare</u>	

STANDARD DATA

UNIVERSITY OF KANSAS, Lawrence, 1966

State: Kansas

Stream: Shosh Creek (1 mi. west of St. Line) Station No. 12, 1966

Location: Sec. 32, T. 34 N., R. 22 E., Cherokee Co., Kansas

Water Temp. (F.) 58° Air Temp. (F.) 75° Turbidity (ft.) 2-3

Rate of Flow (cu. ft./min.) swift Type of drainage: rock-oak-hill

Habitat: limestone bottom, little vegetation in water

Specimens taken

ORDER	FAMILY	SPECIES	REMARKS
CYPRINODONTIFORMES	Cyprinodontidae	<i>Gambusia affinis holbrooki</i>	
		<i>Leuciscus leuciscus</i>	
		<i>Leuciscus xanthurus</i>	
		<i>Leuciscus niloticus</i>	
		<i>Leuciscus niloticus</i>	
		<i>Leuciscus niloticus</i>	
CYPRINODONTIFORMES	Cyprinodontidae	<i>Poecilia reticulata</i>	
		<i>Stethostoma labialis</i>	

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Shoal Creek - Reddings Mill Station Station No. - Date March 19, 1966Location Sec. 6, T. 35 S., R. 33 W., Newton Co., MissouriWater Temp.(F.) 58° Air Temp.(F.) 75° Turbidity(ft.) 2-3Rate of Flow(cu. ft./min.) swift Type of Drainage rocky, wooded hillsHabitat limestone bottom, little vegetation in water; much leaves on bottom

Specimens Taken

ORDER	FAMILY	Species	Remarks
CYPRINIFORMES	Catostomidae	<u>Moxostoma erythrum</u>	
	Cyprinidae	<u>Notemigonus crysoleucas</u>	
		<u>Notropis umbratilis</u>	
	Ictaluridae	<u>Notropis lutrensis</u>	
		<u>Ictaluris melas</u>	
PERCIFORMES	Centrarchidae	<u>Lepomis megalotus</u>	
		<u>Lepomis humilis</u>	
		<u>Lepomis macrochirus</u>	
		<u>Pomoxis nigromaculatus</u>	
		<u>Pomoxis annularis</u>	

CONTINUED DATA

INSTRUMENTAL DATA, Spring, 1966

Site: Shoal Creek - Rabbits Hill Station Station No. - Date: March 12, 1966

Location: Sec. 6, T. 35 S., R. 33 W., Newton Co., Missouri

Water Temp. (F.) 58° Air Temp. (F.) 72° Turbidity (Ft.) 2-3

Rate of Flow (cu. ft./min.) 500 Type of Bottom: rocky, wooded hills

Habitat: Limestone bottom, little vegetation in water; much leaves on bottom

Specimens Taken

ORDER	FAMILY	SPECIES	REMARKS
CYPRINIDAE	Gambusia	Gambusia affinis holbrooki	
	Gambusia	Gambusia affinis holbrooki	
	Gambusia	Gambusia affinis holbrooki	
	Gambusia	Gambusia affinis holbrooki	
	Gambusia	Gambusia affinis holbrooki	
CYPRINIDAE	Gambusia	Gambusia affinis holbrooki	
	Gambusia	Gambusia affinis holbrooki	
	Gambusia	Gambusia affinis holbrooki	
	Gambusia	Gambusia affinis holbrooki	
	Gambusia	Gambusia affinis holbrooki	

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Shoal Creek - Tipton Ford Station Station No. - Date March 19, 1966Location Sec. 15, T. 26 N., R. 32 W., Newton Co., MissouriWater Temp.(F.) 58° Air Temp.(F.) 75° Turbidity(ft.) 2-3Rate of Flow(cu. ft./min.) swift Type of Drainage rocky-wooded hillsHabitat limestone bottom, little vegetation in water; a few leaves on bottom

Specimens Taken

ORDER	FAMILY	Species	Remarks
CYPRINIFORMES	Catostomidae	<u>Hypentelium nigricans</u>	
	Cyprinidae	<u>Hybopsis biguttata</u>	
		<u>Notropis pilsbryi</u>	
		<u>Notropis cornutus</u>	
		<u>Notropis buchanani</u>	
		<u>Campostoma anomalum</u>	
CYPRINODONTIFORMES	Poeciliidae	<u>Gambusia affinis</u>	
PERCIFORMES	Centrarchidae	<u>Micropterus dolomieu</u>	
		<u>Ambloplites rupestris</u>	
	Percidae	<u>Etheostoma nigrum</u>	
		<u>Etheostoma spectabile</u>	
		<u>Etheostoma flabellare</u>	
	Cottidae	<u>Cottus carolinae</u>	

Spokane taken

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Shoal Creek - Low Water Bridge Station Station No.   - Date March 22, 1966Location Sec. 36, T. 34 S., R. 33 W., Newton Co., MissouriWater Temp.(F.) 58° Air Temp.(F.) 75° Turbidity(ft.) 2-3Rate of Flow(cu. ft./min.) swift Type of Drainage typical ozarkianHabitat limestone bottom., much aquatic vegetation

Specimens Taken

ORDER	FAMILY	Species	Remarks
CYPRINIFORMES	Cyprinidae	<u>Notropis boops</u> <u>Notropis camurus</u>	
PERCIFORMES	Centrarchidae	<u>Micropterus dolomieu</u>	
	Percidae	<u>Lepomis microlophus</u> <u>Etheostoma zonale</u> <u>Etheostoma blenniodes</u>	

INSTRUCTIONS

NOVEMBER 22, 1964

Location Sec. 36, T. 35 N., R. 33 W., Newton Co., Missouri

Water Temp. (°C) 20.0 (°F) 68.0

Date of Flow (m. l. rain) 11/11/64

Type of drainage typical of area

Habitat: limestone bottom, much aquatic vegetation

Specimens Taken

ORDER FAMILY SPECIES Remarks

	Pisces	Cyprinidae	<u>Notropis bairdi</u>	
			<u>Notropis cornutus</u>	
			<u>Notropis heterodon</u>	
			<u>Notropis heterodon</u>	
	Pisces	Cyprinidae	<u>Notropis heterodon</u>	
			<u>Notropis heterodon</u>	
			<u>Notropis heterodon</u>	
			<u>Notropis heterodon</u>	
	Pisces	Cyprinidae	<u>Notropis heterodon</u>	
			<u>Notropis heterodon</u>	
			<u>Notropis heterodon</u>	
			<u>Notropis heterodon</u>	

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Shoal Creek - Grand Falls Station Station No. - Date March 26, 1966

Location Sec. 35, T. 34 S., R. 34 W., Newton Co., Missouri

Water Temp.(F.) 58° Air Temp.(F.) 65° Turbidity(ft.) 2-3

Rate of Flow(cu. ft./min.) rapid to very swift Type of Drainage typical ozarkian

Habitat moderatly heavy aquatic vegetation, very rocky, rocks large; current over falls drops 6-10 feet, good darter habitat

Specimens Taken

ORDER	FAMILY	Species	Remarks
CYPRINIFORMES	Cyprinidae	<u>Semotilus atromaculatus</u> <u>Notropis pilsbryi</u> <u>Notropis camurus</u> <u>Notropis lutrensis</u> <u>Campostoma anomalum</u>	
PERCIFORMES	Centrarchidae	<u>MICROPTERUS PUNCTULATUS</u> <u>Lepomis macrochirus</u>	
	Percidae	<u>Lepomis humilis</u> <u>Percina copelandi</u> <u>Etheostoma stigmaeum</u> <u>Etheostoma blenniodes</u> <u>Etheostoma punctulatum</u>	
PERCIFORMES	Centrarchidae	<u>Chaenobryttus gluosus</u>	April 6, 1966

Date: March 23, 1966
 Location: 35.3° N, 122.3° W, Point Barrow, Alaska
 Time: 10:00 AM
 State of sky: Partly cloudy to very light
 Visibility: 10 miles
 Wind: 10-15 mph, light breeze
 Temperature: 10-15°F
 Humidity: 60-70%
 Barometric pressure: 30.0-30.5 in. Hg.
 Remarks: Moderate heavy sea state, very rough, rocks large; current over
 10-15 knots, 5-10 feet, good water habitat

SPECIES	FAMILY	SPECIES	SPECIES
CYPIDIDAE	Cypididae	Cypididae	Cypididae
CYPIDIDAE	Cypididae	Cypididae	Cypididae
CYPIDIDAE	Cypididae	Cypididae	Cypididae
CYPIDIDAE	Cypididae	Cypididae	Cypididae
CYPIDIDAE	Cypididae	Cypididae	Cypididae
CYPIDIDAE	Cypididae	Cypididae	Cypididae

April 6, 1966

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Dry Wood Creek - Lone Star School Station No. 11 Date March 31, 1966Location Sec. 29, T. ~~28~~^{33N.} S., R. ~~25~~^{32W} E., Barton Co., MissouriWater Temp.(F.) 63° Air Temp.(F.) 80° Turbidity(ft.) 1-2Rate of Flow(cu. ft./min.) slow to moderately fast Type of Drainage rocky, wooded hillsHabitat heavy leaf cover on bottom, (large) limestones on bottom; heavy sulfur compounds in water and salts on rocks on banks. probably polluted- no fresh waterSpecimens Taken mollusks in area.

ORDER	FAMILY	Species	Remarks
CYPRINIFORMES	Catostomidae	<u>Catostomus commersoni</u>	
		<u>Cyprinus carpio</u>	
	Cyprinidae	<u>Semotilus atromaculatus</u>	
		<u>Notropis umbratilis</u>	
		<u>Notropis boops</u>	
		<u>Notropis topeka</u>	
		<u>Campostoma anomalum</u>	
		<u>Noturus exilis</u>	
	Ictaluridae		
MUGILIFORMES	Atherinidae	<u>Labidesthes sicculus</u>	
PERCIFORMES	Centrarchidae	<u>Micropterus salmoides</u>	
		<u>Lepomis cyanellus</u>	
		<u>Lepomis megalotus</u>	
		<u>Lepomis humilis</u>	
		<u>Lepomis macrochirus</u>	
		<u>Pomoxis annularis</u>	
	Percidae	<u>Etheostoma spectabile</u>	

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Neosho River Station No. - Date April 10, 1966
Location low water bridge at south edge of Miami, Okla. city park
Water Temp.(F.) 58° Air Temp.(F.) 65° Turbidity(ft.) 1-2 Rate of Flow mod. swift
Type of Drainage populated area, flat, rocky drainage
Habitat river bed very rocky, some organic debree in water
Specimens Taken

ORDER	Family	Species	Remarks
PERCIFORMES	Serranidae	<u>Roccus chrysops</u>	taken on hook and line using minnow for bait male, not yet spawned.

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Dry Wood Creek Station No. 12 & 13 Date April 14, 1966
 Location Sec. 9, T. 33 N., T. 32 W., Barton Co., Missouri (#12=west, #13=east)
 Water Temp.(F.) 50° Air Temp.(F.) 65° Turbidity(ft.) 1-2
 Rate of Flow(cu. ft./min.) slow Type of Drainage rocky wooded hills and
flat farm land on one
 Habitat bottom covered by flat sandstones and clay; much brush in water side
 Specimens Taken

ORDER	FAMILY	Species	Remarks
CYPRINIFORMES	Catostomidae	<u>Catostomus commersoni</u>	
		<u>Notemigonus crysoleucas</u>	
	Cyprinidae	<u>Semotilus atromaculatus</u>	
		<u>Notropis umbratilis</u>	
		<u>Notropis lutrensis</u>	
		<u>Notropis topeka</u>	
		<u>Campostoma anomalum</u>	
		<u>Ictaluris melas</u>	
		<u>Noturus exilis</u>	
	Ictaluridae		
CYPRINODONTIFORMES	Cyprinodontidae	<u>Fundulus notatus</u>	
MUGILIFORMES	Atherinidae	<u>Labidesthes sicculus</u>	
PERCIFORMES	Centrarchidae	<u>Micropterus salmoides</u>	
		<u>Lepomis cyanellus</u>	
		<u>Lepomis megalotus</u>	
		<u>Lepomis humilis</u>	
		<u>Pomoxis annularis</u>	
	Percidae	<u>Etheostoma spectabile</u>	

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Neosho River Station No. - Date April 16, 1966Location Sec. 29, T. 30 S., R. 20 E., Neosho Co., KansasWater Temp.(F.) 54° Air Temp.(F.) 65° Turbidity(ft.) 0-1 Rate of Flow swiftType of Drainage slightly wooded hills and farm landHabitat limestone bottom, some silt and debree on bottom and in water

Specimens Taken

ORDER	Family	Species	Remarks
CYPRINIFORMES	Cyprinidae	<u>Notropis lutrensis</u>	
		<u>Pimephales notatus</u>	
		<u>Campostoma anomalum</u>	
	Ictaluridae	<u>Ictaluris punctatus</u>	
PERCIFORMES	Centrarchidae	<u>Lepomis humilis</u>	
	Percidae	<u>Percina phoxocephala</u>	

Nebraska River, Dec. 10, 1933

Dec. 10, 1933, Nebraska Co., Kansas

3-10-33

slightly wooded hills and farm land

limestone bottom, some silt and debris on bottom and in water

Cyrtopoda luteipes
Cyrtopoda nivalis
Cyrtopoda luteipes

Cyrtopoda luteipes

Cyrtopoda luteipes

Cyrtopoda luteipes

Cyrtopoda

Cyrtopoda

Cyrtopoda

Cyrtopoda

ICHTHYOLOGY 333, Spring, 1966

Location 1 $\frac{1}{4}$ miles southwest of New Strawn, Coffey Co., Kansas

Water Temp.(F.) ? Air Temp.(F.) ? Turbidity(ft.) ? Rate of Flow ?

Type of Drainage_____?

Habitat ?

Specimens Taken

ORDER	Family	Species	Remarks
PERCIFORMES	Sciaenidae	<u>Aplodinotus grunniens</u>	

1938, 12 April - 1938, 12 April - 1938, 12 April

1938, 12 April - 1938, 12 April - 1938, 12 April

1938, 12 April - 1938, 12 April - 1938, 12 April

1938, 12 April - 1938, 12 April - 1938, 12 April

1938, 12 April - 1938, 12 April - 1938, 12 April

1938, 12 April - 1938, 12 April - 1938, 12 April

1938, 12 April - 1938, 12 April - 1938, 12 April

1938, 12 April - 1938, 12 April - 1938, 12 April

1938, 12 April - 1938, 12 April - 1938, 12 April

1938, 12 April - 1938, 12 April - 1938, 12 April

1938, 12 April - 1938, 12 April - 1938, 12 April

1938, 12 April - 1938, 12 April - 1938, 12 April

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Spring River Station No. - Date May 5, 1966
 Location Sec. 19, T. 34 S., R. 25 E., Cherokee Co., Kansas
 Water Temp.(F.) 58° Air Temp.(F.) 75° Turbidity(ft.) 1-2 Rate of Flow moderately
swift
 Type of Drainage wooded hills; drainage somewhat rocky, flat bottom land
 Habitat very rocky, with large boulders in water, quite a bit of aquatic
debris in water
 Specimens Taken

ORDER	Family	Species	Remarks
CYPRINIFORMES	Catostomidae*	<u>Catostomus commersoni</u>	
	Cyprinidae	<u>Notropis rubellus</u> <u>Notropis umbratilis</u> <u>Notropis pilsbryi</u> <u>Notropis boops</u> <u>Notropis lutrensis</u> <u>Notropis buechanani</u> <u>Chrosomus erythrogaster</u> <u>Pimephales notatus</u> <u>Pimephales vigilax</u>	
CYPRINODONTIFORMES	Cyprinodontidae	<u>Fundulus notatus</u>	
	Poeciliidae	<u>Gambusia affinis</u>	
MUGILIFORMES	Atherinidae	<u>Labidesthes sicculus</u>	
PERCIFORMES	Centrarchidae	<u>Micropterus punctulatus</u> <u>Micropterus salmoides</u> <u>Lepomis cyanellus</u> <u>Lepomis megalotus</u> <u>Lepomis humilis</u> <u>Lepomis macrochirus</u> <u>Pomoxis annularis</u>	
	Percidae	<u>Percina caprodes</u> <u>Percina copelandi</u> <u>Etheostoma spectabile</u>	
			*(additions)
CYPRINIFORMES	Catostomidae	<u>Ictiobus bubalus</u> <u>Moxostoma erythrum</u>	

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

0001, 2, 1000

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Dry Wood Creek Station No. - Date May 12, 1966Location Sec. 9, T. 34 N., R. ³²~~33~~ W., Vernon Co., MissouriWater Temp.(F.) 50° Air Temp.(F.) 52° Turbidity(ft.) 0 Rate of Flow mod. swiftType of Drainage flat, sparsely wooded farm land; water backed up into roadside ditches, and reached a considerable depthHabitat (hard to determine due to flood conditions) mostly mud and silt bottom with few rocks; much plant debris in water at this time;Specimens Taken many land organisms in water due to (?) recent storm, especially Insecta and Arachnida, (mostly Fam. Lycosidae.)

ORDER	Family	Species	Remarks
CYPRINIFORMES	Cyprinidae	<u>Notropis umbratilis</u> <u>Notropis lutrensis</u>	
PERCIFORMES	Centrarchidae	<u>Lepomis humilis</u> <u>Lepomis macrochirus</u> <u>Lepomis megalotus</u>	

...and ... ..
... ..
... ..
... ..
... ..
... ..
... ..
... ..
... ..
... ..

airline   airline
airline   airline

<u>airman</u>	airman
<u>airman</u>	airman
<u>airman</u>	airman
<u>airman</u>	airman

sci intro

substantive

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Crawford Co. St. Lake # 2 (Farlington) Station No. - Date August 10, 1966Location Sec. 5, 6, & 7; T. 28 S., R 23 E., Crawford Co., KansasWater Temp.(F.) 81 Air Temp.(F.) 80 Turbidity(ft.) 3.75 Rate of Flow 0Type of Drainage Wooded hills and flat lands, mostly wooded on perimeter of lakeHabitat Clear, man-made impoundment (1934-CCC); little vegetation except along shore-lineCollectors: Hartmann, Tripplett, and Gash

Specimens Taken

ORDER	Family	Species	Remarks
CYPRINIFORMES	Ictaluridae	<u>Ictalurus nebulosis</u>	Taken by electro-fishing Total no. taken: 9; SL ranges from _____ mm to _____ mm.

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Elk City Reservoir Station No. - Date August 16, 1966Location Water Temp.(F.) Air Temp.(F.) Turbidity(ft.) Rate of Flow Type of Drainage Habitat

Specimens Taken Collectors: Hartmann and Tripplet.

ORDER	Family	Species	Remarks
Lepisosteiformes	Lepisosteidae	<u>Lepisosteus osseus</u>	
Cypriniformes	Catostomidae	<u>Ictiobus bubalus</u>	
Perciformes	Percidae	<u>Stizostedion vitreum</u>	
	<u>Cyprinidae</u>	<u>Cyprinus carpio</u>	

1. The purpose of this study is to determine the effect of the treatment on the response of the subjects.

2. The subjects of this study are the patients who have been treated with the drug.

3. The results of the study are as follows:

4. The data were analyzed by the following methods:

5. The results of the analysis are as follows:

6. The results of the analysis are as follows:

7. The results of the analysis are as follows:

8. The results of the analysis are as follows:

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Otter Creek Station No. 2 Date June 28, 1966Location SW $\frac{1}{4}$ of NE $\frac{1}{4}$ of Sec. 16, T. 33 S., R. 8 E., Cowley Co., KansasWater Temp.(F.) - Air Temp.(F.) - Turbidity(ft.) - Rate of Flow -Type of Drainage Flint hills (Chautauqua hills extention).Habitat Clear, smooth rock bottom.

Specimens Taken

ORDER	Family	Species	Remarks
Cypriniformes	Ictaluridae	<u>Noturus flavus</u> <u>Noturus nocturnus</u>	SL: 74 mm SL: 57 mm

1979012115

ICHTHYOLOGY 333, Spring, 1966

Location

Type of Drainage Wooded hills

Habitat mud bottom - few rocks & vegetation

ORDER	Family	Species	Remarks
CYPRINIFORMES	ICTALURIDAE	<u>Pylodictis</u> <u>divaris</u>	1 of 2 taken
			1967

[illegible]

DATE: 11/11/2011 TIME: 11:11 AM (11/11/2011) 11:11 AM

SECRET

And that's it.

10/10/1944

09/06/01

201-103

2000

Journal of Interpersonal Violence

ICHTHYOLOGY 333, Spring, 1966

Location OTTOWA Co, OKLAHOMA

Water Temp. (F.) — Air Temp. (F.) — Turbidity (ft.) — Rate of Flow —

Type of Drainage _____

Habitat Collected by C. OLIVER

ORDER	Family	Species	Remarks
CYPRINIFORMES	CIATOSTOMIDAE	<u>MINYTREMA</u> <u>MELANOPS</u>	SL=120mm

UNITED STATES DEPARTMENT OF AGRICULTURE

Report of the Director of the Bureau of Plant Industry
on the results of the investigation of the
insects and diseases of the cotton plant
in the State of Texas, during the year
1907.

State	County	Name of Grower	Crop
Texas	Dallas	John A. Smith	Cotton
		Wm. H. Jones	Cotton
		Geo. W. Brown	Cotton
		Chas. E. White	Cotton
Texas	Harris	John A. Smith	Cotton
		Wm. H. Jones	Cotton
		Geo. W. Brown	Cotton
		Chas. E. White	Cotton
Texas	Tarrant	John A. Smith	Cotton
		Wm. H. Jones	Cotton
		Geo. W. Brown	Cotton
		Chas. E. White	Cotton
Texas	Cotton	John A. Smith	Cotton
		Wm. H. Jones	Cotton
		Geo. W. Brown	Cotton
		Chas. E. White	Cotton
Texas	Dallas	John A. Smith	Cotton
		Wm. H. Jones	Cotton
		Geo. W. Brown	Cotton
		Chas. E. White	Cotton
Texas	Harris	John A. Smith	Cotton
		Wm. H. Jones	Cotton
		Geo. W. Brown	Cotton
		Chas. E. White	Cotton
Texas	Tarrant	John A. Smith	Cotton
		Wm. H. Jones	Cotton
		Geo. W. Brown	Cotton
		Chas. E. White	Cotton
Texas	Cotton	John A. Smith	Cotton
		Wm. H. Jones	Cotton
		Geo. W. Brown	Cotton
		Chas. E. White	Cotton

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Brazos River Station No. - Date Aug. 17, 1967Location 15 miles west of Bryan, Texas Brazos County.Water Temp.(F.) - Air Temp.(F.) - Turbidity(ft.) - Rate of Flow -Type of Drainage Habitat Rocky bottom, rate of flow only moderate; period of drought.

Specimens Taken

ORDER	Family	Species	No.	Remarks
Cypriniformes	Cyprinidae	<u>Hybopsis aestivalis</u>	2	SL range 21-29 mm
		<u>Pimephales vigilax</u>	3	35-50 mm
		<u>Notropis dorsalis</u> ?	1	35 mm
		<u>Notropis potteri</u>	15	27-45 mm
		<u>Notropis lutrensis</u>	14	21-36 mm
	Ictaluridae	<u>Ictalurus punctatus</u>	2	161-236mm
		<u>Ictalurus furcatus</u>	2	174-196mm
Collection Note: Minnows were collected by seining a shallow (0.5 m) riffle; catfishes were collected by hook-and-line method.				

100-443887-100

10-100-0-11

[illegible][illegible][illegible]

_____

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466

172

100-443887-100

1000

1970-1971 1972-1973 1974-1975 1976-1977 1978-1979 1980-1981 1982-1983 1984-1985 1986-1987 1988-1989 1990-1991 1992-1993 1994-1995 1996-1997 1998-1999 2000-2001 2002-2003 2004-2005 2006-2007 2008-2009 2010-2011 2012-2013 2014-2015 2016-2017 2018-2019 2020-2021 2022-2023 2024-2025 2026-2027 2028-2029 2030-2031 2032-2033 2034-2035 2036-2037 2038-2039 2040-2041 2042-2043 2044-2045 2046-2047 2048-2049 2050-2051 2052-2053 2054-2055 2056-2057 2058-2059 2060-2061 2062-2063 2064-2065 2066-2067 2068-2069 2070-2071 2072-2073 2074-2075 2076-2077 2078-2079 2080-2081 2082-2083 2084-2085 2086-2087 2088-2089 2090-2091 2092-2093 2094-2095 2096-2097 2098-2099 2100-2101 2102-2103 2104-2105 2106-2107 2108-2109 2110-2111 2112-2113 2114-2115 2116-2117 2118-2119 2120-2121 2122-2123 2124-2125 2126-2127 2128-2129 2130-2131 2132-2133 2134-2135 2136-2137 2138-2139 2140-2141 2142-2143 2144-2145 2146-2147 2148-2149 2150-2151 2152-2153 2154-2155 2156-2157 2158-2159 2160-2161 2162-2163 2164-2165 2166-2167 2168-2169 2170-2171 2172-2173 2174-2175 2176-2177 2178-2179 2180-2181 2182-2183 2184-2185 2186-2187 2188-2189 2190-2191 2192-2193 2194-2195 2196-2197 2198-2199 2200-2201 2202-2203 2204-2205 2206-2207 2208-2209 2210-2211 2212-2213 2214-2215 2216-2217 2218-2219 2220-2221 2222-2223 2224-2225 2226-2227 2228-2229 2230-2231 2232-2233 2234-2235 2236-2237 2238-2239 2240-2241 2242-2243 2244-2245 2246-2247 2248-2249 2250-2251 2252-2253 2254-2255 2256-2257 2258-2259 2260-2261 2262-2263 2264-2265 2266-2267 2268-2269 2270-2271 2272-2273 2274-2275 2276-2277 2278-2279 2280-2281 2282-2283 2284-2285 2286-2287 2288-2289 2290-2291 2292-2293 2294-2295 2296-2297 2298-2299 2300-2301 2302-2303 2304-2305 2306-2307 2308-2309 2310-2311 2312-2313 2314-2315 2316-2317 2318-2319 2320-2321 2322-2323 2324-2325 2326-2327 2328-2329 2330-2331 2332-2333 2334-2335 2336-2337 2338-2339 2340-2341 2342-2343 2344-2345 2346-2347 2348-2349 2350-2351 2352-2353 2354-2355 2356-2357 2358-2359 2360-2361 2362-2363 2364-2365 2366-2367 2368-2369 2370-2371 2372-2373 2374-2375 2376-2377 2378-2379 2380-2381 2382-2383 2384-2385 2386-2387 2388-2389 2390-2391 2392-2393 2394-2395 2396-2397 2398-2399 2400-2401 2402-2403 2404-2405 2406-2407 2408-2409 2410-2411 2412-2413 2414-2415 2416-2417 2418-2419 2420-2421 2422-2423 2424-2425 2426-2427 2428-2429 2430-2431 2432-2433 2434-2435 2436-2437 2438-2439 2440-2441 2442-2443 2444-2445 2446-2447 2448-2449 2450-2451 2452-2453 2454-2455 2456-2457 2458-2459 2460-2461 2462-2463 2464-2465 2466-2467 2468-2469 2470-2471 2472-2473 2474-2475 2476-2477 2478-2479 2480-2481 2482-2483 2484-2485 2486-2487 2488-2489 2490-2491 2492-2493 2494-2495 2496-2497 2498-2499 2500-2501 2502-2503 2504-2505 2506-2507 2508-2509 2510-2511 2512-2513 2514-2515 2516-2517 2518-2519 2520-2521 2522-2523 2524-2525 2526-2527 2528-2529 2530-2531 2532-2533 2534-2535 2536-2537 2538-2539 2540-2541 2542-2543 2544-2545 2546-2547 2548-2549 2550-2551 2552-2553 2554-2555 2556-2557 2558-2559 2560-2561 2562-2563 2564-2565 2566-2567 2568-2569 2570-2571 2572-2573 2574-2575 2576-2577 2578-2579 2580-2581 2582-2583 2584-2585 2586-2587 2588-2589 2590-2591 2592-2593 2594-2595 2596-2597 2598-2599 2600-2601 2602-2603 2604-2605 2606-2607 2608-2609 2610-2611 2612-2613 2614-2615 2616-2617 2618-2619 2620-2621 2622-2623 2624-2625 2626-2627 2628-2629 2630-2631 2632-2633 2634-2635 2636-2637 2638-2639 2640-2641 2642-2643 2644-2645 2646-2647 2648-2649 2650-2651 2652-2653 2654-2655 2656-2657 2658-2659 2660-2661 2662-2663 2664-2665 2666-2667 2668-2669 2670-2671 2672-2673 2674-2675 2676-2677 2678-2679 2680-2681 2682-2683 2684-2685 2686-2687 2688-2689 2690-2691 2692-2693 2694-2695 2696-2697 2698-2699 2700-2701 2702-2703 2704-2705 2706-2707 2708-2709 2710-2711 2712-2713 2714-2715 2716-2717 2718-2719 2720-2721 2722-2723 2724-2725 2726-2727 2728-2729 2730-2731 2732-2733 2734-2735 2736-2737 2738-2739 2740-2741 2742-2743 2744-2745 2746-2747 2748-2749 2750-2751 2752-2753 2754-2755 2756-2757 2758-2759 2760-2761 2762-2763 2764-2765 2766-2767 2768-2769 2770-2771 2772-2773 2774-2775 2776-2777 2778-2779 2780-2781 2782-2783 2784-2785 2786-2787 2788

1. The first step in the process is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the problem.

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466

[illegible][illegible][illegible]

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Navasota River Bottom (slough) Station No. - Date Aug. 14, 1967Location Robertson Co., Texas known locally as "Lake Wilburn".Water Temp.(F.) - Air Temp.(F.) - Turbidity(ft.) - Rate of Flow -Type of Drainage Extremely muddy isolated hole.Habitat Pasture-land drainage, but periphery of lake with dead trees.

Specimens Taken

ORDER	Family	Species	Remarks
Amiiformes	Amiidae	<u>Amia calva</u>	1 taken SL _____ mm male, with conspicuous dorsal spot on tail.
Collection Note: Collected by seining.			
+ Same trip - <u>Elanone zonale</u>			

1. The first part of the report (1-10) is a general introduction to the subject of the report. It contains a brief history of the subject and a statement of the purpose of the report.

2. The second part of the report (11-20) is a detailed description of the methods used in the study. It includes a description of the subjects, the apparatus, and the procedures used.

3. The third part of the report (21-30) is a presentation of the results of the study. It includes a description of the data collected and a discussion of the findings.

4. The fourth part of the report (31-40) is a conclusion and a summary of the results. It includes a statement of the conclusions drawn from the study and a summary of the findings.

Introduction	Methods	Results	Conclusions
The first part of the report (1-10) is a general introduction to the subject of the report. It contains a brief history of the subject and a statement of the purpose of the report.	The second part of the report (11-20) is a detailed description of the methods used in the study. It includes a description of the subjects, the apparatus, and the procedures used.	The third part of the report (21-30) is a presentation of the results of the study. It includes a description of the data collected and a discussion of the findings.	The fourth part of the report (31-40) is a conclusion and a summary of the results. It includes a statement of the conclusions drawn from the study and a summary of the findings.

ICHTHYOLOGY 333, Spring, 1966

Specimens Taken

ORDER	Family	Species	Remarks
Anguilliformes	Anguillidae	<u>Anguila</u> <u>rostrata</u>	1972

ICHTHYOLOGY 333, Spring, 1960

Station No. _____ Date _____

Location _____

Water Temp. (F.) _____ Air Temp. (F.) _____ Visibility (ft.) _____ Rate of Flow _____

Type of Drainage _____

Habitat _____

Specimens Taken _____

ORDER	Family	Species	Remarks

COLLECTION DATA

ICHTHYOLOGY 333, Spring, 1966

Area Stony Creek Station No. 6 Date 16 Apr. 1972Location Dauphin Co., PennsylvaniaWater Temp.(F.) 52 Air Temp.(F.) 70 Turbidity(ft.) - Rate of Flow -Type of Drainage -Habitat -

Specimens Taken

ORDER	Family	Species	Remarks
Clupeiformes	Esocidae	Esox niger E. americanus	Caught by angler -Sized Mar. 72.
Cypriniformes	Cyprinidae	<u>Semotilus</u> <u>corporeus</u>	- many in Stony Ck. Collections.

THE UNIVERSITY OF CHICAGO

SECRET

This image shows a single sheet of white paper with horizontal blue or grey ruling lines. A solid vertical line runs down the right side of the page, creating a margin. The paper appears slightly aged or off-white. There are some faint, dark specks scattered across the surface, possibly dust or minor imperfections in the paper or scanning process. No text or other markings are present on the page.