

Pittsburg State University

Pittsburg State University Digital Commons

Posters

Research Colloquium 2019

4-1-2019

Nanosheets of CuCo_2O_4 as a High-Performance Electrocatalyst in Urea Oxidation

Camila Zequine
Pittsburg State University

Ram K. Gupta
Pittsburg State University

Pawan K. Kahol
Pittsburg State University

Follow this and additional works at: https://digitalcommons.pittstate.edu/posters_2019

 Part of the [Energy Systems Commons](#)

Recommended Citation

Zequine, Camila; Gupta, Ram K.; and Kahol, Pawan K., "Nanosheets of CuCo_2O_4 as a High-Performance Electrocatalyst in Urea Oxidation" (2019). *Posters*. 7.
https://digitalcommons.pittstate.edu/posters_2019/7

This Article is brought to you for free and open access by the Research Colloquium 2019 at Pittsburg State University Digital Commons. It has been accepted for inclusion in Posters by an authorized administrator of Pittsburg State University Digital Commons. For more information, please contact lfthompson@pittstate.edu.




Nanosheets of CuCo_2O_4 as a High-Performance Electrocatalyst in Urea Oxidation





Camila Zequine, Pawan K. Kahol, Ram K. Gupta
Pittsburg State University, Pittsburg, KS, United States

Introduction



- Development of new materials capable of generating green energy with high efficiency and low cost.
- Urea is an abundant waste generated in agricultural land which could be used as a source to generate hydrogen.
- Water splitting occurs at 1.23 V to generate hydrogen, while urea oxidation could provide hydrogen at a lower potential of 0.37 V.
- In this work, it was synthesized nanostructured CuCo_2O_4 as an electrocatalyst for urea oxidation to generate hydrogen as a green fuel.



Experimental




Results and discussion




(a) XRD pattern and (b) SEM image of the CuCo_2O_4 .


Structure composed of nanosheets in different orientations and a smooth surface.



(a) Cyclic voltammetry curves in 1 M KOH with and without 0.33 M urea, and (b) corresponding Tafel slopes.





CuCo_2O_4 demanded a potential of **1.36 V**, which is much lower than that without urea.




(a) Polarization curves for CuCo_2O_4 in 1 M KOH without and with 0.33 M urea, (b) Chronopotentiometric curve for CuCo_2O_4 .

Electrolyzer requires a cell voltage of **1.45 V** in urea solution which is 260 mV less than the cell voltage required without urea.



(a) LSV curves of the CuCo_2O_4 electrode toward HER in 1.0 M KOH with 0.33 M urea, (b) corresponding Tafel plots.



CuCo_2O_4 required a potential of **168 mV** to generate 10 mA/cm^2 .

Conclusion and Future work

✓ CuCo_2O_4 in urea oxidation: a solution to eliminate the urea from groundwater systems and environment that are harmful for people; the H_2 production with less energy consumption and low-costs, which helps to solve the crisis energy.

Future work: Make the process even more efficient and develop new multifunctional materials for energy storage and generation.

Acknowledgement

Thanks to Polymer Chemistry Program and Kansas Polymer Research for providing research facilities.