

Pittsburg State University

Pittsburg State University Digital Commons

Posters

Research Colloquium 2019

4-1-2019

Prophylactic Use of LMWH vs UFH in PE Patients

Kyleigh Grieshaber

Pittsburg State University

Barbara McClaskey

Pittsburg State University

Follow this and additional works at: https://digitalcommons.pittstate.edu/posters_2019



Part of the [Cardiovascular Diseases Commons](#), and the [Therapeutics Commons](#)

Recommended Citation

Grieshaber, Kyleigh and McClaskey, Barbara, "Prophylactic Use of LMWH vs UFH in PE Patients" (2019). *Posters*. 22.

https://digitalcommons.pittstate.edu/posters_2019/22

This Article is brought to you for free and open access by the Research Colloquium 2019 at Pittsburg State University Digital Commons. It has been accepted for inclusion in Posters by an authorized administrator of Pittsburg State University Digital Commons. For more information, please contact lfthompson@pittstate.edu.

Prophylactic Use of LMWH vs UFH in PE Patients

Kyleigh Grieshaber, Senior BSN Student

Pittsburg State University, Irene Ransom Bradley School of Nursing

Research Colloquium 2019

Purpose

The purpose of this review of literature was to compare the efficacy, safety, and cost of LMWH with UFH and make decisions about selecting the best prophylactic therapy for PE patients.

PICOT Statement

- **Population:** Patients effected with PE
- **Intervention:** Prophylactic use of LMWH
- **Comparison:** Prophylactic use of UFH
- **Outcome:** decrease in the morbidity and mortality of PE
- **Timeline:** During time of prophylaxis of PE

Interventions

- Unfractionated Heparin (UFH)
 - A Standard dose of UFH consists of an IV bolus of 80U/kg followed by a continuous infusion of 18U/kg/hr.
- Low Molecular Weight Heparin (LMWH)
 - There are many different compounds of LMWH, but the most common ones used are Enoxaparin Sodium, Dalteparin Sodium, and Tinzaparin Sodium. Below, you can see the typical dosing for each.

- Dalteparin sodium 200 units/kg s.c. once daily or 100 units/kg s.c. every 12 hours
- Enoxaparin sodium 1 mg/kg s.c. every 12 hours or 1.5 mg/kg s.c. once daily
- Tinzaparin sodium 175 units/kg s.c. once daily

- LMWH dosage is determined by weight, therefore it can be administered in the out-patient setting
- Both LMWH and UFH require 5-7 initial days of warfarin therapy, with a therapeutic INR goal of 2-3 and aPTT goal of 1.5-2.5x control value. Once these lab values are reached, the patient is on P.O therapy for 3-6 months, depending on the level of risk.

Summary

- UFH and LMWH both decrease the risk of PE with no significant difference in prophylaxis
- LMWH decreases the risk of major hemorrhage by 52% compared to UFH.
- Shorter hospital stays are associated with the use of LMWH because home therapy is an option
- The overall cost of LMWH is less than UFH that could potentially save the U.S. \$250 million a year.

Background

- Pulmonary Embolism (PE) is the third leading cause of cardiovascular death in the United States with at least 950,000 deaths annually.
- PE is the most preventable cause of unexpected deaths in hospitalized patients, however, it is the most common.
- 50-80% of these cases are asymptomatic, with 70% of these cases going undetected until it's too late.
- Virchow's Triad identifies the three major underlying factors that contribute to PE
 - Stasis (changes in blood flow pattern)
 - Injury (changes in blood vessel wall)
 - Hypercoagulability (change in consistency of blood)
- Risk Factors include:
 - Age > 50
 - History of:
 - Varicose veins
 - Myocardial Infarction
 - Cancer
 - Atrial fibrillation
 - Ischemic stroke
 - Diabetes mellitus
- Signs and Symptoms
 - Redness
 - Swelling
 - Rash
 - Hypotension
 - Dyspnea
 - Tachycardia
 - Chest discomfort
 - Hypoxemia
 - Respiratory arrest
 - death

Conclusion

- Overall, the use of LMWH is proven to be more beneficial than the use of UFH in PE prophylaxis.
- There are several advantages including a decrease in the risk of heparin induced thrombocytopenia and hospital length of stay.
- Because LMWH is an effective home therapy due to its consistent effects, it eliminates the need for intense lab monitoring of aPTT and INR values.

Gaps in the Literature

- Larger, more diverse sample size was needed in the article discussing the new JFK Risk Assessment tool.
- Some studies suggested different time frames for long term anticoagulant use, regardless of the use of UFH or LMWH.
- Dosage recommendations for UFH and LMWH fluctuated.
- Overall, the population size was good, validity and reliability were tested in all research studies use in this review of literature.

References

- Copeland DC, & Gretzer MB. (2006). Deep venous thrombosis: evidence-based prevention strategies. *Contemporary Urology*, 18(5), 48-54.
- Dipaola CA. (2008). Patient safety first. Preventing deep vein thrombosis: a perioperative nursing imperative. *AORN Journal*, 88(2), 283-285.
- Elisha, S., Heiner, J., Nagelhout, J., & Gabot, M. (2015). Venous Thromboembolism: New Concepts in Perioperative Management. *AANA Journal*, 83(3), 211-223.
- McCaffrey R, Bishop M, Adonis-Rizzo M, Williamson E, McPherson M, Cruikshank A, ... Lauzon C. (2007). Development and testing of a DVT risk assessment tool: providing evidence of validity and reliability. *Worldviews on Evidence-Based Nursing*, 4(1), 14-20.
- Nadeau C, & Varrone J. (2003). Treat DVT with low molecular weight heparin. *Nurse Practitioner*, 28(10), 22-31.
- Nutescu EA. (2007). Assessing, preventing, and treating venous thromboembolism: evidence-based approaches. *American Journal of Health-System Pharmacy*, 64, S5-13.